arXiv:1612.02144v1 [math.GT] 7 Dec 2016

A q-SERIES IDENTITY VIA THE $\mathfrak{sl}_3$ COLORED JONES POLYNOMIALS FOR THE $(2, 2m)$-TORUS LINK

WATARU YUASA

ABSTRACT. The colored Jones polynomial is a q-polynomial invariant of links colored by irreducible representations of a simple Lie algebra. A q-series called a tail is obtained as the limit of the $\mathfrak{sl}_2$ colored Jones polynomials $\{J_n(K;q)\}_n$ for some link K, for example, an alternating link. For the $\mathfrak{sl}_3$ colored Jones polynomials, the existence of a tail is unknown. We give two explicit formulas of the tail of the $\mathfrak{sl}_3$ colored Jones polynomials colored by $(n, 0)$ for the $(2, 2m)$-torus link. These two expressions of the tail provide an identity of q-series. This is a knot-theoretical generalization of the Andrews-Gordon identities for the Ramanujan false theta function.

1. INTRODUCTION

The Rogers-Ramanujan identities were first discovered and proved by Rogers [19]. After that, these identities appeared in Ramanujan's first letter to Hardy without proof. To this day, the Rogers-Ramanujan identities have been generalized by many people and have been proved in many ways. One of these generalizations is the Andrews-Gordon identities.

Theorem 1.1 (The Andrews-Gordon identities for the Ramanujan theta function [1])**.**

$$f(-q^{2m}, -q) = (q;q)_\infty \sum_{k_m \leq \cdots \leq k_2 \leq k_1} \frac{q^{\sum_{j=1}^{m-1} k_j(k_j+1)}}{\prod_{j=1}^{m-1}(q;q)_{k_j-k_{j+1}}},$$

where $m > 0$ and $f(a,b) = \sum_{i=0}^{\infty} a^{\frac{i(i+1)}{2}} b^{\frac{i(i-1)}{2}} + \sum_{i=1}^{\infty} a^{\frac{i(i-1)}{2}} b^{\frac{i(i+1)}{2}}$ is Ramanujan's general theta function (see, for example, [2]).

In knot theory, the Andrews-Gordon identities appear through the ($\mathfrak{sl}_2$) colored Jones polynomial. The colored Jones polynomial is a q-polynomial invariant of knots and links. Dasbach and Lin showed the stability of some coefficients of the colored Jones polynomials for alternating knots in [6, 7]. They conjectured that the k-th coefficient of the $n+1$ dimensional colored Jones polynomial $J_n(K;q)$ for an alternating knot K coincides up to sign for $n \geq k$. This suggests the existence of q-series $\sum_{k=0}^{\infty} a_k q^k$ for an alternating link K called the *tail*. The coefficient a_k is given by the k-th coefficient of $J_n(K;q)$ with $n \geq k$. Armond [3] proved the existence of tails for adequate links, in particular for alternating knots. Independently, for alternating knots, Garoufalidis and Lê [8] showed a more general stability of coefficients of $J_n(K;q)$. In [5], Armond and Dasbach showed two explicit formulas for the tails of the colored Jones polynomial for the $(2, 2m+1)$-torus knot. These two formulas conclude the left- and right-hand side of the Andrews-Gordon identities for the Ramanujan theta function. They used the formula for the colored Jones polynomial of the $(2, 2m+1)$-torus knot in [16] and obtained by the method of Armond [4]. Hajij [10] also showed the Andrews-Gordon identities for the Ramanujan theta function and the Ramanujan false theta function using two expressions of the tail of the $(2, 2m+1)$-torus knot and the $(2, 2m)$-torus link, respectively.

Theorem 1.2 (The Andrews-Gordon identities for the Ramanujan false theta function [10])**.**

$$\Psi(q^{2m-1},q)=(q;q)_\infty \sum_{k_{m-1}\leq\cdots\leq k_2\leq k_1} \frac{q^{\sum_{j=1}^{m-1}k_j(k_j+1)}}{(q;q)_{k_{m-1}}^2\prod_{j=1}^{m-2}(q;q)_{k_j-k_{j+1}}},$$

where $m>1$ and $\Psi(a,b)=\sum_{i=0}^{\infty}a^{\frac{i(i+1)}{2}}b^{\frac{i(i-1)}{2}}-\sum_{i=0}^{\infty}a^{\frac{i(i-1)}{2}}b^{\frac{i(i+1)}{2}}$ is Ramanujan's general false theta function (see, for example, [15]).

He used formulas for the colored Jones polynomial obtained by using a colored trivalent graph representation of the Kauffman bracket skein elements (see, for example, [12]) and the Kauffman bracket bubble skein expansion formula in [9].

The main subject of this paper is the tail of the colored $\mathfrak{sl}_3$ Jones polynomials for the $(2,2m)$-torus link. The existence of tails of the $\mathfrak{sl}_3$ colored Jones polynomials for any class of knots and links is unknown. Moreover, there is no example of an explicit formula for a tail of the $\mathfrak{sl}_3$ colored Jones polynomial. We will give two explicit formulas of "the $\mathfrak{sl}_3$ tail" for the $(2,2m)$-torus link. As the result, the following q-series identity holds.

Theorem 4.3.

$$\sum_{i=0}^{\infty}q^{-2i}q^{m(i^2+2i)}\frac{(1-q^{i+1})^3(1+q^{i+1})}{1-q}=(q)_\infty\sum_{0\leq k_m\leq\cdots\leq k_2\leq k_1}\frac{q^{-2k_m}q^{\sum_{j=1}^{m}(k_i^2+2k_i)}}{(q)_{k_m}^2(q)_{k_1-k_2}\cdots(q)_{k_{m-1}-k_m}}.$$


The right-hand side of the above identity is obtained from the full twist formula in [21]. The left-hand side is obtained by using of colored trivalent graph presentation of A_2 webs.

The paper is organized as follows. We first review definitions and formulas related to the A_2 web space in section 2. The A_2 web space is a generalization of the Kauffman bracket skein module. In section 3, we introduce a method to represent some types of A_2 webs using colored trivalent graphs. Furthermore, we give values of some θ-graphs and quantum $6j$ symbols. As an application, we explicitly give the $\mathfrak{sl}_3$ colored Jones polynomial for a 2-bridge link. In section 4, we derive two q-series, that is "the $\mathfrak{sl}_3$ tails", by using explicit formulas for the $\mathfrak{sl}_3$ colored Jones polynomial for the $(2,2m)$-torus link in section 3 and in [21].

2. The A_2 web space and some formulas

The skein theory has been developed with a quantum representation of Lie algebra $\mathfrak{g}$. If $\mathfrak{g}=\mathfrak{sl}_2$, the skein theory consists of the Kauffman bracket skein module, which is called the Temperley-Lieb algebra and the A_1 web space, and the Kauffman bracket. Kuperberg constructed the skein theory for Lie algebras of rank 2 in [13, 14]. In this section, we review definitions of the A_2 web space, the A_2 bracket and the A_2 clasp.

Let $\varepsilon=(\varepsilon_1,\varepsilon_2,\ldots,\varepsilon_m)$ be an m-tuple of signs $+$ or $-$. Let D_ε denote the unit disk with signed marked points $\{\exp(2\pi\sqrt{-1}/m)^{j-1}\mid j=1,2,\ldots,m\}$ on its boundary. The sign of $\exp(2\pi\sqrt{-1}/m)^{j-1}$ is given by ε_j for $j=1,2,\ldots,m$. A *bipartite uni-trivalent graph* G is a directed graph such that each vertex is either trivalent or univalent and the vertices are divided into the sinks and the sources. A sink (resp. source) is a vertex such that all edges adjoining to the vertex point into (resp. away from) it. A *bipartite trivalent graph* G in D_ε is an embedding of a uni-trivalent graph into D_ε such that any vertex v has the following neighborhoods:







FIGURE 2.1. The Reidemeister moves for tangled trivalent graph diagrams

An A_2 *basis web* is the boundary-fixing isotopy class of a bipartite trivalent graph G in D_ε, where any internal face of $D_\varepsilon \setminus G$ has at least six sides. Let us denote B_ε the set of A_2 basis webs in D_ε. For example, $B_{(+,-,+,-,+,-)}$ has the following A_2 basis webs:



The A_2 *web space* W_ε is the $\mathbb{Q}(q^{\frac{1}{6}})$-vector space spanned by B_ε. A *tangled trivalent graph diagram* in D_ε is an immersed bipartite uni-trivalent graph in D_ε whose intersection points are only transverse double points of edges with crossing data  or  . Tangled trivalent graph diagrams G and G' are regularly isotopic if G is obtained from G' by a finite sequence of boundary-fixing isotopies and Reidemeister moves, see Figure 2.1, with some direction of edges.

Tangled trivalent graphs in D_ε are regular isotopy classes of tangled trivalent graph diagrams in D_ε. We denote T_ε the set of tangled trivalent graphs in D_ε.

Definition 2.1 (The A_2 bracket [14])**.** We define a $\mathbb{Q}(q^{\frac{1}{6}})$-linear map $\langle \cdot \rangle_3 \colon \mathbb{Q}(q^{\frac{1}{6}})T_\varepsilon \to W_\varepsilon$ by the following.

- $\left\langle \right\rangle_3 = q^{\frac{1}{3}} \left\langle \right\rangle_3 - q^{-\frac{1}{6}} \left\langle \right\rangle_3,$
- $\left\langle \right\rangle_3 = q^{-\frac{1}{3}} \left\langle \right\rangle_3 - q^{\frac{1}{6}} \left\langle \right\rangle_3,$
- $\left\langle \right\rangle_3 = \left\langle \right\rangle_3 + \left\langle \right\rangle_3,$
- $\left\langle \right\rangle_3 = [2] \left\langle \right\rangle_3,$
- $\left\langle G \sqcup \right\rangle_3 = [3] \langle G \rangle_3,$


where $[n] = \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}$ is a quantum integer.

We remark that this map is invariant under the Reidemeister moves for tangled trivalent graphs.

We next consider the A_2 web space $W_{n^+ +n^-} = W_{(+,+,\dots,+,-,-,\dots,-)}$ whose first n marked points are decorated with $+$ and next n marked points are decorated with $-$. We define A_2 clasps  $\in W_{n^+ +n^-}$ inductively by the following.

Definition 2.2. (The A_2 clasps)

 $\in W_{1^+ +1^-}$

(2.1)  $\in W_{n^+ +n^-}$


A_2 clasps have the following properties.

Lemma 2.3 (Properties of A_2 clasps)**.** For any positive integer n,


-  ,
- 

We also define the A_2 clasp of type (n, m) according to Ohtsuki and Yamada [17].

Definition 2.4 (the A_2 clasp of type (n, m))**.**



Lemma 2.5 (Property of A_2 clasps of type (m, n))**.**



We review some formulas for clasped A_2 web spaces in [21]. We define a q-Pochhammer symbol as

$$(q;q)_k = \prod_{l=1}^{k}(1-q^l).$$

We abbreviate it as $(q)_k$. A q-binomial coefficient is defined by

$$\binom{n}{k}_q = \frac{(q;q)_n}{(q;q)_k(q;q)_{n-k}}$$

for $k \leq n$. If $k > n$, we define it by 0. We also define a q-multinomial coefficient as

$$\binom{n}{n_1, n_2, \dots, n_m}_q = \frac{(q)_n}{(q)_{n_1}(q)_{n_2}\cdots(q)_{n_m}},$$

where $n_1, n_2, \dots, n_m$ are non-negative integers such that $n_1 + n_2 + \cdots + n_m = n$.

Theorem 2.6 (The m full twists formula [21])**.**

$$\left\langle \quad \right\rangle_3 = q^{-\frac{2m}{3}(n^2+3n)} \sum_{0\le k_m\le\cdots\le k_1\le n} q^{n-k_m} q^{\sum_{i=1}^{m}(k_i^2+2k_i)}$$

$$\times \frac{(q)_n}{(q)_{k_m}} \binom{n}{k_1', k_2', \ldots, k_m', k_m}_q \left\langle \quad \right\rangle_3,$$

where k_i, k_i' are integers such that $k_0 = n$, $k_{i+1}' = k_i - k_{i+1}$ for $i = 0, 1, \ldots, m-1$.

Theorem 2.7 (The A_2 bracket bubble skein expansion formula [21])**.**

$$\left\langle \quad \right\rangle_3 = \sum_{t=\max\{k,l\}}^{\min\{k+l,n,m\}} \frac{\begin{bmatrix} n \\ t \end{bmatrix}\begin{bmatrix} m \\ t \end{bmatrix}\begin{bmatrix} t \\ k \end{bmatrix}\begin{bmatrix} t \\ l \end{bmatrix}\begin{bmatrix} n+m-t+2 \\ n+m-k-l+2 \end{bmatrix}}{\begin{bmatrix} n \\ k \end{bmatrix}\begin{bmatrix} m \\ k \end{bmatrix}\begin{bmatrix} n \\ l \end{bmatrix}\begin{bmatrix} m \\ l \end{bmatrix}} \left\langle \quad \right\rangle_3,$$

where $\begin{bmatrix} a \\ b \end{bmatrix} = \frac{[a]!}{[b]![a-b]!}$ for $a \ge b$ and $[a]! = \prod_{i=1}^{a} [i]$.

3. TRIVALENT GRAPHS FOR CLASPED A_2 WEB SPACES

For the Kauffman bracket skein module, a method for representing its elements by colored trivalent graphs is well known. Formulas related to colored trivalent graphs play a very important role in defining the quantum $SU(2)$ invariants for 3-manifolds and the quantum representations for mapping class groups of surfaces, etc. (See, for example, Kauffman and Lins [12], Turaev and Viro [20] and Roberts [18].) In this section, we represent a certain clasped A_2 web space by using colored trivalent graphs and give some formulas.

Definition 3.1. Let n be a non-negative integer. For $0 \le i \le n$, we define a colored trivalent graph by .

We use the following notations:

- $\Delta(m, n) = \left\langle \quad \right\rangle_3$,
- $\theta(n, n, (i, i)) = \left\langle \quad \right\rangle_3$,
- $\mathrm{Tet}\begin{bmatrix} n & n & (j,j) \\ n & n & (i,i) \end{bmatrix} = \left\langle \quad \right\rangle_3$,
- $\begin{Bmatrix} n & n & (j,j) \\ n & n & (i,i) \end{Bmatrix} = \dfrac{\mathrm{Tet}\begin{bmatrix} n & n & (j,j) \\ n & n & (i,i) \end{bmatrix} \Delta(j,j)}{\theta(n,n,(j,j))^2}$,


where m, n are any non-negative integers and $0 \le i, j \le n$.

Lemma 3.2.

(1) $\Delta(i,j) = \frac{[i+1][j+1][i+j+2]}{[2]}$,


(2) $\theta(n,n,(i,i)) = \sum_{k=0}^{i}(-1)^k \frac{\begin{bmatrix} i \\ k \end{bmatrix}^2}{\begin{bmatrix} 2i+1 \\ k \end{bmatrix}} \frac{\Delta(n,0)^2}{\Delta(n-i+k,0)} = \frac{\begin{bmatrix} n+i+2 \\ 2i+2 \end{bmatrix}}{\begin{bmatrix} n \\ i \end{bmatrix}^2}\Delta(i,i)$,

(3) $\mathrm{Tet}\begin{bmatrix} n & n & (j,j) \\ n & n & (i,i) \end{bmatrix} = \sum_{k=0}^{i}(-1)^k \frac{\begin{bmatrix} i \\ k \end{bmatrix}^2 \begin{bmatrix} n-j \\ i-k \end{bmatrix} \begin{bmatrix} n+j+2 \\ i-k \end{bmatrix}}{\begin{bmatrix} 2i+1 \\ k \end{bmatrix} \begin{bmatrix} n \\ i-k \end{bmatrix}^2}\theta(n,n,(j,j))$.

Proof. We only show (2) and (3). The first equation of (2) is obtained by expanding the double-lined edge using Definition 2.4. $\theta(n,n,(i,i))$ is also represented by a colored graph $\left\langle$  $\right\rangle_3$. This clasped A_2 web has a bubble skein element colored by $n-i$ and we use Theorem 2.7. Thus, we obtain the second equation of (2). The proof of (3) is obtained by using Definition 2.4 and Theorem 2.7. We expand a double-lined edge colored by s. Then, the clasped A_2 web appearing in each term has two bubble skein elements. Finally, $\theta(n,n,(j,j))$ is obtained by expanding one of these bubble skein elements. □

Lemma 3.3.

$$\left\langle \quad \right\rangle_3 = \delta_{ij}\frac{\theta(n,n,(i,i))}{\Delta(i,i)}\left\langle \quad \right\rangle_3,$$




where $\overset{i}{=\!=}$ denotes  and δ_{ij} is the Kronecker delta function.

Proof. Theorem 2.7 and Lemma 2.5 show the value is zero if $i \neq j$. If $i = j$, then the coefficient is obtained by closing double-lined edges of both sides. □

Proposition 3.4 (Recoupling Theorem)**.**

$$\left\langle \quad \right\rangle_3 = \sum_{j=0}^{n}\begin{Bmatrix} n & n & (j,j) \\ n & n & (i,i) \end{Bmatrix}\left\langle \quad \right\rangle_3$$




Proof. We can prove this proposition in the same way as the recoupling theory of Temperley-Lieb algebra, see [12, Chapter 7], using Definition 2.4 and Lemma 3.3. □

We give a formula of the $\mathfrak{sl}_3$ colored Jones polynomial $J^{\mathfrak{sl}_3}_{(n,0)}([2a_1, 2a_2, \ldots, 2a_l]\,;q)$ for a 2-bridge link $[2a_1, 2a_2, \ldots, 2a_l]$ using colored trivalent graphs. In this paper we define the $\mathfrak{sl}_3$ colored Jones polynomial for framed links obtained by a link diagram with the blackboard framing.


Definition 3.5.

$$J^{\mathfrak{sl}_3}_{(n,0)}([2a_1, 2a_2, \ldots, 2a_l]\,;q)$$

$$= \begin{cases} \left\langle \quad \right\rangle_3 \Big/ \Delta(n,0) & \text{if } l \text{ is odd,} \\ \left\langle \quad \right\rangle_3 \Big/ \Delta(n,0) & \text{if } l \text{ is even,} \end{cases}$$




where, $a_1, a_2, \ldots, a_l$ are non-zero integers and




Theorem 3.6.

$$J^{\mathfrak{sl}_3}_{(n,0)}([2a_1, 2a_2, \ldots, 2a_l]\,; q)$$
$$= \sum_{0\le i_1, i_2 \ldots i_l \le n} \frac{\Delta(i_1,i_1)}{\Delta(n,0)} \frac{\theta(n,n,(i_l,i_l))}{\theta(n,n,(i_1,i_1))} q^{-\frac{2}{3}(n^2+3n)(\sum_{k=1}^{l} a_k)} q^{\sum_{k=1}^{l} a_k(i_k^2+2i_k)}$$
$$\times \prod_{k=1}^{l-1} \begin{Bmatrix} n & n & (i_{k+1}, i_{k+1}) \\ n & n & (i_k, i_k) \end{Bmatrix}.$$

Lemma 3.7.




Proof. It is shown by using the following formula [21, Lemma 3.16]:




and Theorem 2.6 with $m = 1$. □

Proof of Theorem 3.6.

(3.1)


This equation is derived from Proposition 3.4, Lemma 3.7 and $\begin{Bmatrix} n & n & (i_k,i_k) \\ n & n & (0,0) \end{Bmatrix} = \frac{\Delta(i_k,i_k)}{\theta(n,n,(i_k,i_k))}$. By Proposition 3.4 and Lemma 3.3,

(3.2)


First, we apply (3.1) to boxed $2a_k$ for all $0 \le k \le l$ of the two bridge link diagram. Next, we repeatedly apply (3.2) to the leftmost doubled edge. Finally, we obtain $\theta(n, n, (i_l, i_l))$. □

4. The tail of the $\mathfrak{sl}_3$ colored Jones polynomial for $T(2,2m)$

We discuss a stability of coefficients of the $\mathfrak{sl}_3$ colored Jones polynomial for the torus link $T(2,2m)$ in this section for positive integer m. Two explicit formulas of $J^{\mathfrak{sl}_3}_{(n,0)}(T(2,2m))$ are obtained from Theorem 3.6 and [21, Theorem 5.7] because $T(2,2m)$ has a presentation $[2m]$ as a 2-bridge link. We write these two formulas below. Let m be a positive integer.

From Theorem 3.6,

$$\begin{aligned}\psi^{(m)}_n(q) = J^{\mathfrak{sl}_3}_{(n,0)}(T(2,2m)) &= q^{-\frac{2m}{3}(n^2+3n)}\sum_{i=0}^{n}\frac{\Delta(i,i)}{\Delta(n,0)}q^{m(i^2+2i)}\\ &= q^{-\frac{2m}{3}(n^2+3n)+n}\sum_{i=0}^{n}q^{-2i}q^{m(i^2+2i)}\frac{(1-q^{i+1})^3(1+q^{i+1})}{(1-q)(1-q^{n+1})(1-q^{n+2})}.\end{aligned}$$

From [21, Theorem 5.7],

$$\begin{aligned}g^{(m)}_n(q) &= J^{\mathfrak{sl}_3}_{(n,0)}(T(2,2m))\\ &= q^{-\frac{2m}{3}(n^2+3n)+n}\sum_{0\le k_m\le\cdots\le k_2\le k_1\le n} q^{-2k_m}q^{\sum_{j=1}^{m}(k_i^2+2k_i)}\\ &\quad\times\frac{(q)_n^2}{(q)_{k_m}^2(q)_{n-k_1}(q)_{k_1-k_2}\cdots(q)_{k_{m-1}-k_m}}\frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{n-k_m+1})(1-q^{n-k_m+2})}.\end{aligned}$$

Definition 4.1. Suppose $f(q), f_n(q)\in\mathbb{Z}[[q]]$ for $n\ge 1$. *The limit of* $\{f_n(q)\}_n$ *is* $f(q)$, denoted $\lim_{n\to\infty}f_n(q)=f(q)$, means that $f_n(q)=f(q)$ in $\mathbb{Z}[[q]]/q^{n+1}\mathbb{Z}[[q]]$ for all n.

We can see that the minimum degree of $J^{\mathfrak{sl}_3}_{(n,0)}(T(2,2m))$ is $q^{-\frac{2m}{3}(n^2+3n)+n}$ and consider the limit of $\Psi^{(m)}_n(q)=q^{\frac{2m}{3}(n^2+3n)-n}\psi_n(q)$ and $G^{(m)}_n(q)=q^{\frac{2m}{3}(n^2+3n)-n}g_n(q)$. The following lemma ensures the existence of the limit.

Lemma 4.2. $\Psi^{(m)}_n(q)=\Psi^{(m)}_{n+1}(q)$ in $\mathbb{Z}[[q]]/q^{n+1}\mathbb{Z}[[q]]$.

Proof. It is clear from the expression of $\Psi^{(m)}_{n+1}(q)$. □

We explicitly give the limit of $\{\Psi^{(m)}_n(q)\}_n$ and $\{G^{(m)}_n\}_n$. First, we can easily obtain the following:

$$\lim_{n\to\infty}\Psi^{(m)}_n(q)=\sum_{i=0}^{\infty}q^{-2i}q^{m(i^2+2i)}\frac{(1-q^{i+1})^3(1+q^{i+1})}{1-q}. \tag{4.1}$$

Next, we consider the limit of $\{G^{(m)}_n\}_n$. In $\mathbb{Z}[[q]]/q^{n+1}\mathbb{Z}[[q]]$,

$$\begin{aligned}\frac{q^{k_m^2}}{1-q^{n-k_m+1}} &= q^{k_m}(1+q^{n-k_m+1}+q^{2(n-k_m+1)}+\dots)\\ &= q^{k_m^2}+q^{n+1+k_m^2-k_m}+(\text{ higher-order tems })\\ &= q^{k_m^2}\end{aligned}$$

because $k_m^2-k_m\ge 0$. In a similar way, we obtain $\frac{q^{k_m^2}}{1-q^{n-k_m+2}}=q^{k_m^2}$, $q^{k_1^2+2k_1}\frac{(q)_n}{(q)_{n-k_1}}=q^{k_1^2+2k_1}$. Form the above, we conclude that the limit of $\{G^{(m)}_n\}_n$ is

$$\lim_{n\to\infty}G^{(m)}_n(q)=(q)_\infty\sum_{0\le k_m\le\cdots\le k_2\le k_1}\frac{q^{-2k_m}q^{\sum_{j=1}^{m}(k_i^2+2k_i)}}{(q)_{k_m}^2(q)_{k_1-k_2}\cdots(q)_{k_{m-1}-k_m}} \tag{4.2}$$

Consequently, we have the following identity of q-series.

Theorem 4.3.

$$\sum_{i=0}^{\infty} q^{-2i} q^{m(i^2+2i)} \frac{(1-q^{i+1})^3(1+q^{i+1})}{1-q} = (q)_\infty \sum_{0 \le k_m \le \cdots \le k_2 \le k_1} \frac{q^{-2k_m} q^{\sum_{j=1}^{m}(k_i^2+2k_i)}}{(q)_{k_m}^2 (q)_{k_1-k_2} \cdots (q)_{k_{m-1}-k_m}}.$$

The above identity is a knot-theoretical generalization of the Andrews-Gordon identities for the Ramanujan false theta function.

Question 4.4. Does the left-hand side of the identity of Theorem 4.3 have a representation as a well-known function like the Ramanujan false theta function?

We compute the expansion of q-series $\Psi^{(m)}(q) = \lim_{n\to\infty} \Psi_n^{(m)}(q)$ up to order 150 by Mathematica 11.0 [11] in Table 1. It is conjectured that the collection of coefficients of the expansion of $\Psi^{(m)}(q)$ except for zeros is $\{1, -1, -1, 1, 1, 1, -1, -1, -1, -1, 1, 1, 1, 1, 1, \dots\}$ for $m > 0$. Sequences of non-zero coefficients and zero coefficients appear in the expansion of $\Psi^{(m)}(q)$ alternatively. The n-th sequence of non-zero coefficients has length $4n$ and the n-th sequence of zero coefficients length $(2n+1)(m-2)$ for $m > 0$.

Acknowledgment. The author would like to express his gratitude to his adviser, Professor Hisaaki Endo, for his encouragement.

REFERENCES

1. George E. Andrews, *An analytic generalization of the Rogers-Ramanujan identities for odd moduli*, Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 4082–4085. MR 0351985
2. George E. Andrews and Bruce C. Berndt, *Ramanujan's lost notebook. Part I*, Springer, New York, 2005. MR 2135178
3. Cody Armond, *The head and tail conjecture for alternating knots*, Algebr. Geom. Topol. **13** (2013), no. 5, 2809–2826. MR 3116304
4. ———, *Walks along braids and the colored Jones polynomial*, J. Knot Theory Ramifications **23** (2014), no. 2, 1450007, 15. MR 3197051
5. Cody Armond and Oliver T. Dasbach, *Rogers-ramanujan type identities and the head and tail of the colored jones polynomial*, arXiv:1106.3948 (2011).
6. Oliver T. Dasbach and Xiao-Song Lin, *On the head and the tail of the colored Jones polynomial*, Compos. Math. **142** (2006), no. 5, 1332–1342. MR 2264669
7. ———, *A volumish theorem for the Jones polynomial of alternating knots*, Pacific J. Math. **231** (2007), no. 2, 279–291. MR 2346497
8. Stavros Garoufalidis and Thang T. Q. Lê, *Nahm sums, stability and the colored Jones polynomial*, Res. Math. Sci. **2** (2015), Art. 1, 55. MR 3375651
9. Mustafa Hajij, *The Bubble skein element and applications*, J. Knot Theory Ramifications **23** (2014), no. 14, 1450076, 30. MR 3312619
10. ———, *The tail of a quantum spin network*, Ramanujan J. **40** (2016), no. 1, 135–176. MR 3485997
11. Wolfram Research Inc., *Mathematica*, version 11.0, Champaign, Illinois (2016).
12. Louis H. Kauffman and Sóstenes L. Lins, *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, Annals of Mathematics Studies, vol. 134, Princeton University Press, Princeton, NJ, 1994. MR 1280463 (95c:57027)
13. Greg Kuperberg, *The quantum G_2 link invariant*, Internat. J. Math. **5** (1994), no. 1, 61–85. MR 1265145
14. ———, *Spiders for rank 2 Lie algebras*, Comm. Math. Phys. **180** (1996), no. 1, 109–151. MR 1403861
15. James McLaughlin, Andrew V. Sills, and Peter Zimmer, *Rogers-Ramanujan computer searches*, J. Symbolic Comput. **44** (2009), no. 8, 1068–1078. MR 2523768
16. H. R. Morton, *The coloured Jones function and Alexander polynomial for torus knots*, Math. Proc. Cambridge Philos. Soc. **117** (1995), no. 1, 129–135. MR 1297899
17. Tomotada Ohtsuki and Shuji Yamada, *Quantum $\mathrm{SU}(3)$ invariant of 3-manifolds via linear skein theory*, J. Knot Theory Ramifications **6** (1997), no. 3, 373–404. MR 1457194
18. Justin Roberts, *Skeins and mapping class groups*, Math. Proc. Cambridge Philos. Soc. **115** (1994), no. 1, 53–77. MR 1253282

m	$\Psi^{(m)}(q)$
1	$1+O\left(q^{151}\right)$
2	$1-q-q^2+q^3+q^4+q^5-q^6-q^7-q^8-q^9+q^{10}+q^{11}+q^{12}+q^{13}+q^{14}-q^{15}$ $-q^{16}-q^{17}-q^{18}-q^{19}-q^{20}+q^{21}+q^{22}+q^{23}+q^{24}+q^{25}+q^{26}+q^{27}-q^{28}-q^{29}$ $-q^{30}-q^{31}-q^{32}-q^{33}-q^{34}-q^{35}+q^{36}+q^{37}+q^{38}+q^{39}+q^{40}+q^{41}+q^{42}+q^{43}$ $+q^{44}-q^{45}-q^{46}-q^{47}-q^{48}-q^{49}-q^{50}-q^{51}-q^{52}-q^{53}-q^{54}+q^{55}+q^{56}+q^{57}$ $+q^{58}+q^{59}+q^{60}+q^{61}+q^{62}+q^{63}+q^{64}+q^{65}-q^{66}-q^{67}-q^{68}-q^{69}-q^{70}-q^{71}$ $-q^{72}-q^{73}-q^{74}-q^{75}-q^{76}-q^{77}+q^{78}+q^{79}+q^{80}+q^{81}+q^{82}+q^{83}+q^{84}+q^{85}$ $+q^{86}+q^{87}+q^{88}+q^{89}+q^{90}-q^{91}-q^{92}-q^{93}-q^{94}-q^{95}-q^{96}-q^{97}-q^{98}-q^{99}$ $-q^{100}-q^{101}-q^{102}-q^{103}-q^{104}+q^{105}+q^{106}+q^{107}+q^{108}+q^{109}+q^{110}+q^{111}$ $+q^{112}+q^{113}+q^{114}+q^{115}+q^{116}+q^{117}+q^{118}+q^{119}-q^{120}-q^{121}-q^{122}-q^{123}$ $-q^{124}-q^{125}-q^{126}-q^{127}-q^{128}-q^{129}-q^{130}-q^{131}-q^{132}-q^{133}-q^{134}-q^{135}$ $+q^{136}+q^{137}+q^{138}+q^{139}+q^{140}+q^{141}+q^{142}+q^{143}+q^{144}+q^{145}+q^{146}+q^{147}$ $+q^{148}+q^{149}+q^{150}+O(q^{151})$
3	$1-q-q^2+q^3+q^7+q^8-q^9-q^{10}-q^{11}-q^{12}+q^{13}+q^{14}+q^{20}+q^{21}+q^{22}-q^{23}$ $-q^{24}-q^{25}-q^{26}-q^{27}-q^{28}+q^{29}+q^{30}+q^{31}+q^{39}+q^{40}+q^{41}+q^{42}-q^{43}-q^{44}$ $-q^{45}-q^{46}-q^{47}-q^{48}-q^{49}-q^{50}+q^{51}+q^{52}+q^{53}+q^{54}+q^{64}+q^{65}+q^{66}+q^{67}$ $+q^{68}-q^{69}-q^{70}-q^{71}-q^{72}-q^{73}-q^{74}-q^{75}-q^{76}-q^{77}-q^{78}+q^{79}+q^{80}+q^{81}$ $+q^{82}+q^{83}+q^{95}+q^{96}+q^{97}+q^{98}+q^{99}+q^{100}-q^{101}-q^{102}-q^{103}-q^{104}-q^{105}$ $-q^{106}-q^{107}-q^{108}-q^{109}-q^{110}-q^{111}-q^{112}+q^{113}+q^{114}+q^{115}+q^{116}+q^{117}$ $+q^{118}+q^{132}+q^{133}+q^{134}+q^{135}+q^{136}+q^{137}+q^{138}-q^{139}-q^{140}-q^{141}-q^{142}$ $-q^{143}-q^{144}-q^{145}-q^{146}-q^{147}-q^{148}-q^{149}-q^{150}+O\left(q^{151}\right)$
4	$1-q-q^2+q^3+q^{10}+q^{11}-q^{12}-q^{13}-q^{14}-q^{15}+q^{16}+q^{17}+q^{28}+q^{29}+q^{30}$ $-q^{31}-q^{32}-q^{33}-q^{34}-q^{35}-q^{36}+q^{37}+q^{38}+q^{39}+q^{54}+q^{55}+q^{56}+q^{57}-q^{58}$ $-q^{59}-q^{60}-q^{61}-q^{62}-q^{63}-q^{64}-q^{65}+q^{66}+q^{67}+q^{68}+q^{69}+q^{88}+q^{89}+q^{90}$ $+q^{91}+q^{92}-q^{93}-q^{94}-q^{95}-q^{96}-q^{97}-q^{98}-q^{99}-q^{100}-q^{101}-q^{102}+q^{103}$ $+q^{104}+q^{105}+q^{106}+q^{107}+q^{130}+q^{131}+q^{132}+q^{133}+q^{134}+q^{135}-q^{136}-q^{137}$ $-q^{138}-q^{139}-q^{140}-q^{141}-q^{142}-q^{143}-q^{144}-q^{145}-q^{146}-q^{147}+q^{148}+q^{149}$ $+q^{150}+O\left(q^{151}\right)$
5	$1-q-q^2+q^3+q^{13}+q^{14}-q^{15}-q^{16}-q^{17}-q^{18}+q^{19}+q^{20}+q^{36}+q^{37}+q^{38}$ $-q^{39}-q^{40}-q^{41}-q^{42}-q^{43}-q^{44}+q^{45}+q^{46}+q^{47}+q^{69}+q^{70}+q^{71}+q^{72}-q^{73}$ $-q^{74}-q^{75}-q^{76}-q^{77}-q^{78}-q^{79}-q^{80}+q^{81}+q^{82}+q^{83}+q^{84}+q^{112}+q^{113}$ $+q^{114}+q^{115}+q^{116}-q^{117}-q^{118}-q^{119}-q^{120}-q^{121}-q^{122}-q^{123}-q^{124}-q^{125}$ $-q^{126}+q^{127}+q^{128}+q^{129}+q^{130}+q^{131}+O\left(q^{151}\right)$
6	$1-q-q^2+q^3+q^{16}+q^{17}-q^{18}-q^{19}-q^{20}-q^{21}+q^{22}+q^{23}+q^{44}+q^{45}+q^{46}$ $-q^{47}-q^{48}-q^{49}-q^{50}-q^{51}-q^{52}+q^{53}+q^{54}+q^{55}+q^{84}+q^{85}+q^{86}+q^{87}-q^{88}$ $-q^{89}-q^{90}-q^{91}-q^{92}-q^{93}-q^{94}-q^{95}+q^{96}+q^{97}+q^{98}+q^{99}+q^{136}+q^{137}$ $+q^{138}+q^{139}+q^{140}-q^{141}-q^{142}-q^{143}-q^{144}-q^{145}-q^{146}-q^{147}-q^{148}-q^{149}$ $-q^{150}+O\left(q^{151}\right)$

TABLE 1. Expansion of $\Psi^{(m)}(q)$ up to order 150 for $m=1,2,\ldots,6$.

19. L. J. Rogers, *Second Memoir on the Expansion of certain Infinite Products*, Proc. London Math. Soc. **S1-25** (1894), no. 1, 318. MR 1576348
20. V. G. Turaev and O. Ya. Viro, *State sum invariants of* 3*-manifolds and quantum* $6j$*-symbols*, Topology **31** (1992), no. 4, 865–902. MR 1191386
21. Wataru Yuasa, *The* $\mathfrak{sl}_3$ *colored Jones polynomials for* 2*-bridge links*, arXiv:1609.07289 (2016).

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1 OOKAYAMA, MEGURO-KU, TOKYO 152-8551, JAPAN
E-mail address: yuasa.w.aa@m.titech.ac.jp